

THE LATEST

**DTN Livestock Open:
Cattle Futures Uneasy
On Cash Market
Pressure** 10-15



**DTN Grain Open: Mixed
on South American
Buying, Export
Interest** 10-15



**DTN Livestock Closing:
Contracts Elevated** 10-14




**ARC, PLC Enrollment
Begins for 2021** 10-14




**DTN Grain Closing:
Soybeans Higher** 10-14



**DTN Cotton Closing:
Cotton Hangs Higher** 10-14




**DTN Livestock Midday:
Lean Hog Market
Boosting** 10-14



**DTN Grain Midday: Corn
and Beans Higher** 10-14



**DTN Cotton Open:
Market Higher,
Challenges
Resistance** 10-14



**Global Markets: Rice –
Brazil Seeks Imports
from Non-Traditional
Suppliers** 10-13



**Global Markets: Wheat – Pasta
Consumption Drives Rebound in**

Soybean Herbicide Injury: Dicamba isn't the Only Culprit

July 4, 2018

By Bob Hartzler and Meaghan Anderson,
Iowa State University Extension Specialists



Photo: Ohio State University

While off-target dicamba injury to soybean has dominated the news the past year, it is important to recognize that dicamba is not the only Group 4 herbicide (HG4) capable of injuring soybean.

These herbicides mimic the activity of indole acetic acid (IAA), a hormone that regulates the activity of numerous genes involved in plant growth. IAA also is referred to as auxin. HG4 products can induce plant responses at lower doses than most other herbicide groups, thus off-target injury has been a problem since their introduction in the 1940's.

This article will discuss some of the problems observed this growing season. All HG4 cause malformed leaves, and distinguishing symptoms between products is difficult (Figure 1).

Timing of symptom development and patterns of injury are important in identifying the source of injury.



Figure 1. Typical dicamba symptoms on soybean.

WEATHER

Local Weather
Radar
U.S. Satellite
24-Hour Surface
F'cast
24-Hour Rainfall Est.
48-Storm Path Accum.
6-10 Day Temp F'cast
6-10 Day Precip.
F'cast
Thunderstorm
Outlook
US Drought Monitor
US Palmer Drought
Index
US Crop Moisture
Index
US Growing Degree
Days
Hurricane Tracks

MARKETS

Cotton
Futures/Options
Cotton Daily
Cotton Weekly
Soybeans
Futures/Options
Corn Futures/Options
Wheat
Futures/Options
Rice Futures/Options
Rice – Milled
Weekly Grain Review
All Commodities

EDITOR'S
PICKS

USDA
ARC, PLC Enrollment
Begins for 2021 10-14




Ag Equipment Sales
Up in Sept. in All Size
Segments 10-13

USDA Announces
Oct. 30 Deadline to
Submit Wildfire,


Global Durum Exports 10-13



Global Markets: Corn – EU-Ukraine Trade Tapers Down 10-13




Global Markets: Oilseeds – U.S. Soybean Futures Rise on Strong Chinese Demand, Limited Brazilian Supplies 10-13



Global Markets: Cotton – China State Reserve Sales Finish on Target 10-13




Corn, Soybeans: Smaller Supply Forecast, as Derecho Harvest Impact Grows 10-13



Ag Equipment Sales Up in Sept. in All Size Segments 10-13



Corn Production: Are 20-Inch Rows the Future? – DTN 10-13



Thompson on Cotton: WASDE Pushes Market Out of 8-Month Trading Range 10-12



Ag Trade: Soybeans, Meat, Corn – What's Affecting the Markets? 10-12



Livestock: Cattle Finishing Net Returns in 2020 10-12



Respiratory Protection During Harvest – Video 10-12



USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster

Soybean Herbicide Injury: Dicamba isn't the Only Culprit – AgFax

Carryover following use in corn

Clopyralid is sold individually as Stinger, but in corn is more commonly used in the premixes Hornet and SureStart. While clopyralid has a much longer half-life than 2,4-D and dicamba, the rates used in corn typically do not carryover at toxic concentrations into the following season.

However, many areas of the state were rain deficient during the 2017 growing season, leading to increased persistence. When clopyralid residues are responsible for damage to soybean, symptoms typically appear by the V1 stage (first trifoliolate). Damage is usually associated with soil type or in streaks related to spray overlaps, rather than field wide injury.

Spread of contaminated hay or planting of soybeans into former pastures

HG4 are commonly used in pastures and hay fields for broadleaf control. Grazon P+D and GrazonNext are popular products that contain picloram and aminopyralid, respectively. Like clopyralid, these two herbicides are relatively persistent.

These herbicides can persist at phytotoxic concentrations in the soil, in forage harvested from treated fields, and in manure of animals consuming forage. Problems may develop when bales of hay from treated fields are placed or spread in fields to be planted to soybean (Figure 2).

Concentrations can be high enough to severely damage, or even kill, emerging soybean. Several instances of this type of injury have been observed this year.




Figure 2. HG4 injury from spreading hay from field treated with picloram on field planted to soybean.

Drift from adjacent fields

Undoubtedly, movement of HG4 from treated areas is the most common source of off-target injury. Off-target injury from use of dicamba in Xtend soybean was a significant problem in Iowa in 2017. Last year we estimated that 150,000 acres of soybean in Iowa were damaged.


However, it is important to recognize that HG4 are used in other areas than soybean. We are aware of numerous situations this season of dicamba used in corn damaging adjacent soybean, and each year HG4 use in roadsides results in damage to sensitive plants.



Hurricane Disaster Assistance Applications 10-12




Biofuels: A Blend of Ethanol, Politics – DTN 10-9



Farm Input Chain Changing, Retailers Going Digital – DTN 10-9

COTTON



West Texas Cotton: Foliar Symptoms Driven by Weather, Not Disease


Louisiana: Assessing Crops in Hurricane Delta's Aftermath

ARC, PLC Enrollment Begins for 2021

DTN Cotton Closing: Cotton Hangs Higher

DTN Cotton Open: Market Higher, Challenges Resistance

GRAIN



DTN Grain Open: Mixed on South

Assistance Applications 10-12



Shurley on Cotton: The Long March to 68 – Now What? 10-12




Weekly Cotton Market Review – USDA 10-9




Biofuels: A Blend of Ethanol, Politics – DTN 10-9



Farm Input Chain Changing, Retailers Going Digital – DTN 10-9



Soybean Market: Bullish Trading – Don't Get Too Excited, Optimistic – DTN 10-9



Midwest Soybeans: Why Is My Crop So Short, What Should I Do About Harvest? – DTN 10-9



Cleveland on Cotton: Fundamentals Remain Weak, 70 Cents Unlikely 10-9




WASDE Cotton: U.S. Production Down Slightly, Ending Stocks Unchanged 10-9



WASDE Oilseeds: U.S. Soybean Production Down on Reduced Harvested Area 10-9



WASDE Rice: Domestic Usage Unchanged, Ending Stocks Raised 10-9



WASDE Coarse Grains: Lower Domestic Corn Production, Usage 10-9

Soybean Herbicide Injury: Dicamba isn't the Only Culprit – AgFax

Dicamba use in Xtend soybean poses a greater risk than when used in corn since it is applied later in the season. Numerous label changes and required training for applicators using dicamba on soybean should help reduce the frequency of off-target damage, but it is too early in the season to determine the impact of these changes.

In 2017, reports of dicamba injury in soybean did not begin until after July 4 in north central Iowa. Numerous reports of injury have been reported in 2018, at this time mostly in southern Iowa.

Mistaken sources of injury

The widespread occurrence of herbicide resistant weeds has resulted in an increase in both the quantity and frequency of herbicide applications in soybean. This increases the likelihood of adverse crop responses – sometimes dicamba gets blamed for damage that it isn't responsible for, and sometimes dicamba injury is blamed on other factors.

The distinct symptom of dicamba is cupping of leaves that emerge after exposure; this injury is typically not noticed for 7-14 days after application due to the time it takes for new leaves to emerge following exposure. The number of leaves affected is determined by the rate of exposure.

All HG4 cause distorted leaf growth, but there can be differences in the type of leaf malformations caused by different herbicides. Frequently 2,4-D results in elongated, strapped leaves in contrast to the cupping normally observed with dicamba (Figure 3).

However, dicamba-type cupping occasionally may be caused by other HG4, and elongated leaflets may develop following dicamba exposure. Thus, it is always critical to identify the source of problem rather than assigning blame based solely on the symptomatology observed.



Figure 3. Elongated leaflet associated with preplant 2,4-D application with sprayer malfunction resulting in excessive rate.

American Buying, Export Interest

Illinois: How Have Acreage Decisions Shifted Over Time?

Iowa: Cool Stored Grain Now

Louisiana: Assessing Crops in Hurricane Delta's Aftermath

ARC, PLC Enrollment Begins for 2021



ARC, PLC Enrollment Begins for 2021

Florida Peanuts: 2020 Rosa County Variety Trial – Video Tour

Florida Weather: La Niña Here to Stay for a While

USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications




ARC, PLC Enrollment Begins for 2021

USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications

Biofuels: A Blend of Ethanol, Politics – DTN

RICE



WASDE Wheat: Higher Domestic Use, Lower Ending Stocks 10-9



Soybeans: Chaff Lining Fields for Palmer Amaranth Management 10-9



Rice Market Update: Louisiana Braces for Hurricane Delta as Industry Digests Harvest Data 10-9



Drought Monitor Weekly: Mostly Cool and Dry, Warmer in the West 10-9



Moving Grain: New Facility Expects to Revive Barge Traffic on Northern Missouri River 10-9



USDA: Peanut Price Highlights 10-9



[More News Items](#)

Send press releases to Ernst@Agfax.com.

[View All Events](#)

Texas: Fall CEU Conference, Robstown, Oct. 15

Texas: Ag Pesticide Waste Collection, Lubbock, Oct. 21

Minnesota: Soil Management Summit, Mankato, Dec. 15-16

Send press releases to Ernst@Agfax.com.

[View All Events](#)

There has been discussion that AMS used with Liberty and other postemergence products may be the source of leaf cupping. AMS has a long history of use as a spray additive, and leaf cupping is not a plant response associated with AMS. The use of group 15 herbicides (Dual, Warrant, Zidua, etc.) has greatly increased in soybean to improve waterhemp control.

These herbicides may cause abnormal development of leaves, but the symptoms do not involve the veinal distortion typical of dicamba or other HG4. The classic symptom of preemergence applications of HG15 is shortening of the midrib on leaflets, resulting in heart-shaped leaflets. Post applications can also cause this symptomology, or they can cause other distortion of leaflets, resulting in irregular margins of leaflets (Figure 4).

Sometimes, distortion of developing leaves can happen with POST applications of the HG14 contact products as well. This distortion is generally accompanied by some contact burn.




Figure 4. Leaf malformations following postemergence application of HG 15 product. Note lack of symmetry in symptoms on leaflets.

Group 4 herbicides are important tools for managing weeds in a variety of situations. While effective tools, their ability to induce plant responses at fractions of label rates requires a higher level of management than other herbicides. When symptoms of HG4 appear it is important to identify the source of the herbicide, and take steps to avoid repeat occurrences in the future.

Tagged [soybeans](#), [herbicide injury](#), [soybean news](#), [Bob Hartzler](#), [auxin herbicides](#), [dicamba injury](#), [Meaghan Anderson](#), [group 4 herbicides](#), [growth regulator herbicides](#)

Source:

<https://crops.extension.iastate.edu/cropnews/2018/07/crop-injury-associated-growth-regulator-herbicides>

Cotton

Grain



Louisiana: Assessing Crops in Hurricane Delta's Aftermath


ARC, PLC Enrollment Begins for 2021

Louisiana: Deja Vu as Hurricane Delta Hits Rice Country

Global Markets: Rice – Brazil Seeks Imports from Non-Traditional Suppliers

USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications

TREE AND VINE



USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications

Drought Monitor Weekly: Mostly Cool and Dry, Warmer in the West


WEATHER



Louisiana: Assessing Crops in Hurricane Delta's Aftermath

Louisiana: Deja Vu as Hurricane Delta Hits Rice Country

Florida Weather: La Niña Here to Stay for




[West Texas Cotton: Foliar Symptoms Driven by Weather, Not Disease](#)

[Louisiana: Assessing Crops in Hurricane Delta's Aftermath](#)

[ARC, PLC Enrollment Begins for 2021](#)

[DTN Cotton Closing: Cotton Hangs Higher](#)

[DTN Cotton Open: Market Higher, Challenges Resistance](#)



[DTN Grain Open: Mixed on South American Buying, Export Interest](#)

[Illinois: How Have Acreage Decisions Shifted Over Time?](#)

[Iowa: Cool Stored Grain Now](#)

[Louisiana: Assessing Crops in Hurricane Delta's Aftermath](#)

[ARC, PLC Enrollment Begins for 2021](#)

a While

Drought Monitor Weekly: Mostly Cool and Dry, Warmer in the West

Peanuts



[ARC, PLC Enrollment Begins for 2021](#)

[Florida Peanuts: 2020 Rosa County Variety Trial – Video Tour](#)

[Florida Weather: La Niña Here to Stay for a While](#)

[USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications](#)

Policy



[ARC, PLC Enrollment Begins for 2021](#)

[USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications](#)

[Biofuels: A Blend of Ethanol, Politics – DTN](#)

Rice

[Louisiana: Assessing Crops in Hurricane Delta's Aftermath](#)
[ARC, PLC Enrollment Begins for 2021](#)

[Louisiana: Deja Vu as Hurricane Delta Hits Rice Country](#)

[Global Markets: Rice – Brazil Seeks Imports from Non-Traditional Suppliers](#)

[USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications](#)

Tree and Vine

[USDA Announces Oct. 30 Deadline to Submit Wildfire, Hurricane Disaster Assistance Applications](#)

[Drought Monitor Weekly: Mostly Cool and Dry, Warmer in the West](#)

Weather

Louisiana: Assessing Crops in
Hurricane Delta's Aftermath

Louisiana: Deja Vu as Hurricane
Delta Hits Rice Country

Florida Weather: La Niña Here
to Stay for a While

Drought Monitor Weekly: Mostly
Cool and Dry, Warmer in the
West

Industry/Press Releases

**Stephen Censky Returns as
CEO of American Soybean
Association** **Discovery Park of America
Moves Closer to Million-
Dollar Goal for New Ag
Innovation Exhibit**

**Corn And Soybean Yields:
Barchart Ups Its U.S.
Projections** **Barchart Offering Free
Commodity Data, Tools To
University Ag Econ
Programs**

**Future Farmers of
America: Record
Membership Numbers,
Diversity** **Soybeans: New South
American Futures Contract
Launches in Sept.**

**U.S. Patents Granted for
Envita Nitrogen-Fixing
Technology**

More Grain Info

AgFax © 2020 All Rights Reserved.